

Lower Bounds on the Algebraic Immunity of Boolean Functions

Hao Chen

Department of Computing and
Information Technology

Fudan University

Shanghai 200433, P.R.China

and

Jianhua Li

Department of Electronic Engineering

Shanghai JiaoTong University

Shanghai 200030, P.R.China

May, 2006

Abstract

From the motivation of algebraic attacks to stream and block ciphers([1,2,7,13,14,15]), the concept of *algebraic immunity* (AI) of a Boolean function was introduced in [21] and studied in [3,5,10,11,17,18,19,20,21]. High algebraic immunity is a necessary condition for resisting algebraic attacks. In this paper, we give some lower bounds on the algebraic immunity of Boolean functions. The results are applied to give lower bounds on the AI of symmetric Boolean functions and rotation symmetric Boolean functions. Some balanced rotation symmetric Boolean functions with their AI near the maximum possible value $\lceil \frac{n}{2} \rceil$ are constructed.

Index Terms— Algebraic attack, Boolean function, algebraic immunity, symmetric Boolean function, rotation symmetric Boolean function

I. Introduction and Preliminaries

A Boolean function of n variable is a mapping $f : F_2^n \rightarrow F_2$, where F_2 is the field of two elements. The weight of a Boolean function $wt(f) = |S_1(f)|$, where $S_1(f) = \{(x_1, \dots, x_n) : f(x_1, \dots, x_n) = 1\}$ and $|\cdot|$ is the cardinality of the set. Any Boolean function has its algebraic normal form (ANF)

$$f(x_1, \dots, x_n) = a_0 + \sum_{i_1 < \dots < i_t} a_{i_1, \dots, i_t} x_{i_1} \cdots x_{i_t}$$

, where $a_0, \dots, a_{i_1, \dots, i_t} \in F_2$. The (algebraic) degree of f is the number of variables in the highest order term in the above ANF. The Boolean function of degree 1 is called affine form. Given a Boolean function f of n variables, a n variable Boolean function g is called its annihilator function if $gf = 0$, or equivalently, g is zero at all points of $S_1(f)$. A Boolean function is called balanced if the number of points in $S_1(f)$, $wt(f) = 2^{n-1}$. The distance of two Boolean functions f and g is $d(f, g) = |S_1(f - h)|$. The nonlinearity of a Boolean function F is defined as $NL(f) = \min_l \{d(f, l)\}$

where l takes over all possible affine forms (see [9]).

Boolean functions are widely used in block and stream ciphers, f.g., in S-boxes, combination generators and filter generators. It is known that Boolean functions used in the practice of cryptography have to satisfy some criteria, f.g., their degrees and nonlinearities etc have to be high (see [9]). Algebraic attack was proposed recently to block and stream ciphers (see [1],[2],[7],[13],[14],[15]). Because of some successful algebraic attacks to several keystream generators, now it is interested to understand the algebraic immunity $AI(f)$ of a Boolean function f , which was introduced in [21]. General properties about algebraic immunity of Boolean functions have been studied in [3],[10],[11],[17],[19],[20],[21]. High algebraic immunity is a necessary condition (but not sufficient) for resisting algebraic attacks. It was proved that the AI of a n variable Boolean function is less than or equal to $\lceil \frac{n}{2} \rceil$ (see [21]). Recently several algorithms for the computation for AI of Boolean functions were given in [4]. If the $AI(f)$ of a Boolean function f is relatively small, the algorithms can be used to determine the $AI(f)$ efficiently. However it is also known that there are Boolean functions of n variables with their AI equal to the maximal possible value $\lceil \frac{n}{2} \rceil$ (see [5],[10],[12],[18]). Thus it is interesting to know more Boolean functions with their AI equal to or near the upper bound $\lceil \frac{n}{2} \rceil$.

A Boolean function is called symmetric if its value is determined by the weight of its input vector. Symmetric Boolean functions have been studied by many authors(see [8] and references there) from the motivation of block and stream ciphers. In software and hardware implementation the symmetric Boolean functions are efficient. Thus it is interested to know the properties of AI of symmetric Boolean functions. In [5], the algebraic immunity of symmetric Boolean functions was thoroughly studied. The AI of elementary symmetric Boolean functions was explicitly determined and some symmetric functions of maximum possible AI have been constructed. Rotation symmetric Boolean functions (RSBF) were introduced and studied in [22] for the purpose of fast hashing. A Boolean function f on F_2^n is called rotation symmetric if $f(x_1, x_2, \dots, x_n) = f(x_n, x_1, \dots, x_{n-1})$ for any $(x_1, x_2, \dots, x_n) \in F_2^n$. The experimental studies of the algebraic immunity of RSBF was initiated in [17]. From the motivation of the possible use of symmetric and rotation symmetric Boolean functions in cryptography, we are interested to have lower bounds on the algebraic immunity of these functions and the construction of these functions with relative high algebraic immunity.

We recall some basic facts about the algebraic immunity of a n variable Boolean function(see [21],[10],[19],[3]).

Definition. Let f be a Boolean function on F_2^n , its algebraic immunity $AI(f)$ is defined to be the smallest number k , such that, there exists one Boolean function g of degree k which is the annihilator function of f or $1 + f$.

Theorem 1 (see [10],[21],[17]). Let f be a n variable Boolean function. Then 1) $AI(f) \leq \lceil \frac{n}{2} \rceil$; 2) $NL(f) \geq 2^{\sum_{i=0}^{AI(f)-2} C_{n-1}^i}$, where C_u^j is the binomial coefficient; 3) If $AI(f) > d$ then $\sum_{i=0}^d C_n^i \leq wt(f) \leq \sum_{i=0}^{n-(d+1)} C_n^i$.

Theorem 2(see [3]). Let f be a Boolean function of n variables. Suppose $wt(f) \geq 2^n - 2^{n-d}$. Then any annihilator of f has its algebraic degree at least d .

We note that Theorem 2 can not be applied directly to *balanced* Boolean functions when lower bounding the AI of Boolean functions. As far as our knowledge, there are quite few *explicitly given* Boolean functions with the maximal possible AI and people do not know much about how to lower bound the algebraic immunity of Boolean functions (see [10],[12],[17],[18]). In this paper we apply Theorem 2 to the restrictions of Boolean functions on some affine subspaces of F_2^n . Thus we present a method to obtain some lower bounds on the algebraic immunity of Boolean functions. In this case, it is possible that the restrictions of the annihilator functions on the affine subspaces are zero. However if the affine subspaces are taken sufficiently many, this consideration leads to some useful results on the lower bound for the AI of Boolean functions.

II. Main Result

The following Theorem 3 is the main result of this paper.

Theorem 3. *If f is a Boolean function on F_2^n and L_1 (respectively L_2) is an affine subspaces with dimension t (respectively s), such that $|S_1(f|_{L_1})| > 2^t - 2^{t-d}$ (respectively $|S_1((1+f)|_{L_2})| > 2^s - 2^{s-d}$). Then*

- 1) *either the annihilator functions of f with minimum possible degree (respectively the annihilator functions of $1+f$ with minimum possible degree) have their degree at least d or;*
- 2) *the annihilator functions of f with minimum possible degree (respectively the annihilator functions of $1+f$ with minimum possible degree) are zero on L_1 (respectively on L_2).*

When Theorem 3 is applied to the balanced Boolean functions and codimension 1 affine subspace we have the following simple conclusion. The proof of Corollary 1 is a direct application of Theorem 3.

Corollary 1. *Let f be a balanced Boolean function on F_2^n and l is an affine form on F_2^n . Suppose $d(f, l) \geq 2^n - 2^{n-d}$. Then we have,*

- 1) *either the algebraic immunity $AI(f)$ is at least d or;*
- 2) *the annihilator functions of f with the minimum possible degree or the annihilator functions of $1+f$ with the minimum possible degree contain l as a factor.*

In section III we can use Theorem 3 to give lower bounds on the algebraic immunity of some symmetric and rotation symmetric Boolean functions by using sufficiently many affine subspaces.

We also have the following result about the Hamming weight of the restrictions of Boolean functions on affine subspaces.

Corollary 2. *Let f be a Boolean function on F_2^n with $AI(f) = d+1$ and L be a affine subspace of F_2^n with codimension r . Then the Hamming weight of f restricted on L satisfies $\sum_{i=0}^{d-r} C_{n-r}^i \leq wt(f|_L) \leq \sum_{i=0}^{n-(d+1)} C_{n-r}^i$.*

When Corollary 2 applied to symmetric Boolean functions we have the following result.

Corollary 3. *Let f be a n variable symmetric Boolean function. Then f can not have the*

maximal possible algebraic immunity $\lceil \frac{n}{2} \rceil$ in the following two cases.

- 1) When n is odd and $wt(x) \geq \lfloor \frac{n}{2} \rfloor$, $f(x)$ is 1 only when $wt(x)$ is odd (or only when $wt(x)$ is even), $f(x)$ can be arbitrary for $wt(x) < \lfloor \frac{n}{2} \rfloor$.
- 2) When n is even and $wt(x) \geq \frac{n}{2} - 1$, $f(x)$ is 1 only when $wt(x)$ is odd (or only when $wt(x)$ is even), $f(x)$ can be arbitrary for $wt(x) < \frac{n}{2} - 1$.

By computing $d(f, l)$, where l is the affine form $x_1 + \dots + x_n$ or $x_1 + \dots + x_n + 1$, and applying Corollary 2, we have the conclusion of Corollary 3 immediately.

Proof of Theorem 3. Let g be an annihilator function of f , that is $gf = 0$. We have $(g|_{L_1})(f|_{L_1}) = 0$. From Theorem 2 $g|_{L_1}$ has its algebraic degree at least d if it is not a zero function. The conclusion is proved.

Proof of Corollary 2. Let l_1, \dots, l_r be r linearly independent affine forms such that L is defined by $l_1 = \dots = l_r = 0$. Considering the Boolean function $f|_L$ as a Boolean function of $n - r$ variables, if its algebraic immunity is smaller $d - r$, we have a Boolean function g' of $n - r$ variables with algebraic degree at most $d - r$ such that $g'(f|_L) = 0$ or $g'((1 + f)|_L) = 0$. Thus the Boolean function $g = (l_1 + 1) \cdots (l_r + 1)g'$ can be think as a Boolean function of n variables of algebraic degree at most d . We have $gf = 0$ or $g(1 + f) = 0$. This is a contradiction. Therefore the algebraic immunity of $f|_L$ is at least $d - r + 1$, we have the conclusion of 1) from the Theorem 1.

III. Lower Bound for AI of Symmetric and Rotation Symmetric Boolean Functions

In this section we use the main result to prove some lower bounds on the algebraic immunity of symmetric and rotation symmetric Boolean functions.

A. Symmetric Boolean Functions

Corollary 4. Let f be a n variable symmetric Boolean function with simplified value vector $v(f) = (v_0(f), \dots, v_i(f), \dots, v_n(f))$, i.e., $f(x) = v_i(f)$ when $wt(x) = i$. Set

$$U = \min\{\sum_{v_i(f)=1, i \leq \lfloor n/2 \rfloor} C_{\lfloor n/2 \rfloor}^i, \sum_{v_i(f)=0, i \geq \lfloor n/2 \rfloor} C_{\lfloor n/2 \rfloor}^{i - \lfloor n/2 \rfloor}\}$$

Suppose $U > 2^{\lfloor n/2 \rfloor} - 2^{\lfloor n/2 \rfloor - d}$. Then $AI(f) \geq d + 1$.

Proof. Let $i_1, \dots, i_{\lfloor \frac{n}{2} \rfloor}$ be arbitrary $\lfloor \frac{n}{2} \rfloor$ indices, L_b be the dimension $\lfloor \frac{n}{2} \rfloor$ subspace of F_2^n defined by $x_{i_1} = \dots = x_{i_{\lfloor \frac{n}{2} \rfloor}} = b$, where $b = 0$ or $b = 1$. If the condition of Corollary 4 is satisfied, $S_1(f|_{L_0}) > 2^{\lfloor n/2 \rfloor} - 2^{\lfloor n/2 \rfloor - d}$ and $S_1((1 + f)|_{L_1}) > 2^{\lfloor n/2 \rfloor} - 2^{\lfloor n/2 \rfloor - d}$. From Theorem 3, either $AI(f) > d$ or the annihilator functions of f or $1 + f$ with minimum possible degree are zero on L_0 and L_1 . This implies that the monomials in the algebraic normal forms f (and $1 + f$) have to contain at least $\lfloor \frac{n}{2} \rfloor$ variables. In the later case $AI(f) = \lfloor \frac{n}{2} \rfloor$. The conclusion is proved.

Example 1. Let f be a 15 variable symmetric Boolean function $f = \sigma_2 + \sigma_4 + \sigma_6 + \sigma_{10} + \sigma_{12} + \sigma_{14}$. Then we have its simplified value vector $v_f = (0, 0, 1, 1, 1, 1, 1, 1, 0, 0, 0, 0, 0, 0, 1, 1)$. Then $U = 246 > 240$ and $AI(f) \geq 5$

Example 2. Let f be a n variable symmetric Boolean function, $I = \{1, \dots, \lfloor \frac{n}{2} \rfloor, n - i\} - \{i\}$ where $i \leq \lfloor \frac{n}{2} \rfloor$, $J = \{\lceil \frac{n}{2} \rceil, \dots, n, i\} - \{n - i\}$. The symmetric Boolean function is defined as follows.

$$\begin{aligned} f(x) &= 1, wt(x) \in I \\ f(x) &= 0, wt(x) \in J \end{aligned}$$

Let t be the smallest positive integer such that $C_{\lceil \frac{n}{2} \rceil}^i + 1 < 2^t$. It is clear $t < i \log_2 n - i$. We have $U > 2^{\lceil \frac{n}{2} \rceil} - 2^t$ and $AI(f) \geq \lceil \frac{n}{2} \rceil - t + 1$. It is obvious that t is asymptotically less than $i \log_2 n$. These Boolean functions have their algebraic immunities asymptotically larger than $n/2 - i \log_2 n + i - 1$.

It is observed from Corollary 4 and Example 2, for a symmetric Boolean function f with the property that most vectors in $S_1(f)$ have their weight less than $\lceil \frac{n}{2} \rceil$ and most vectors in $S_0(f)$ have their weight larger than $\lceil \frac{n}{2} \rceil$, its AI is relatively high. This suggests that these symmetric Boolean functions can be possibly used in stream ciphers, if they satisfy other cryptographic criteria.

B. Rotation Symmetric Boolean Functions

In this subsection we use Theorem 3 to give lower bound for the algebraic immunity of RSBFs.

Example 3. Let f be a rotation symmetric Boolean function of 6 variable

$$\begin{aligned} f = & x_1x_2x_3 + x_2x_3x_4 + x_3x_4x_5 + x_4x_5x_6 + x_5x_6x_1 + x_6x_1x_2 \\ & + x_1x_4 + x_2x_5 + x_3x_6 + x_1x_3x_5 + x_2x_4x_6 + \\ & x_1x_2x_3x_4 + x_2x_3x_4x_5 + x_3x_4x_5x_6 + x_4x_5x_6x_1 + \\ & x_5x_6x_1x_2 + x_6x_1x_2x_3 + x_1x_2x_3x_4 \end{aligned}$$

This is a balanced Boolean function with nonlinearity 24 and $\Delta(f) = 40$, which satisfies $PC(2)$ criteria (see [24]).

We consider two affine subspaces L_1 (respectively L_2) in F_2^6 defined by $x_1 = x_2 = x_3 = 0$ (respectively $x_1 = 1, x_2 = x_3 = 0$). It is easy to check that $S_1((1 + f)|_{L_1})$ has 7 points (in L_1) and $S_1(f|_{L_2})$ has 5 points (in L_2). Thus the annihilator functions of $1 + f$ (respectively, f) have degree at least 2 or are zero on L_1 (respectively L_2). In the later case, the annihilator functions of $1 + f$ (respectively, f) are zero on any rotation transformation of L_1 (respectively, L_2). From this observation, we have $AI(f) \geq 2$.

Example 4. It is clear that each orbit in F_2^n under the circular action $\rho(x_1, x_2, \dots, x_n) = (x_n, x_1, \dots, x_{n-1})$ contains h elements, where h is a factor of n . On the other hand the orbit of a weight i vector in F_2^n under the action of all permutations contains C_n^i elements, which is the union of orbits of circular actions.

From [5] and [8] we know the following *Balanced* symmetric Boolean function f of n (n is odd) variables has the maximal possible AI $\lceil \frac{n}{2} \rceil$.

$$\begin{aligned} f(x) &= 1, wt(x) < \lceil \frac{n}{2} \rceil \\ f(x) &= 0, wt(x) \geq \lceil \frac{n}{2} \rceil \end{aligned}$$

When n is even, the value b in the following definition can be suitably chosen such that it is balanced (in this case the function is not symmetric, however it can be rotation symmetric if b is chosen to be the same on the orbits of circular actions).

$$\begin{aligned} f(x) &= 1, wt(x) < \frac{n}{2} \\ f(x) &= 0, wt(x) < \frac{n}{2} \\ f(x) &= b \in F_2, wt(x) = \frac{n}{2} \end{aligned}$$

If we exchange some orbits under circular actions in the two sets $S_0(f)$ and $S_1(f)$, we get some rotation symmetric Boolean functions and the lower bound on their AI can be proved by applying Theorem 3. Let $H \subset S_0(f)$ and $H' \subset S_1(f)$ be two subsets with the same cardinality, which are the union of orbits under circular actions. Set $X = S_0(f) \cup H' - H$, $X' = S_1(f) \cup H - H'$. Let f' be the Boolean function with $S_0(f') = X$, $S_1(f') = X'$. This is a balanced Boolean function. We have the following result.

Corollary 5. $AI(f') > \lceil \frac{n}{2} \rceil - \lceil \log_2 |H| \rceil$.

When n goes to infinity, we have constructed some balanced rotation symmetric Boolean functions with their algebraic immunity asymptotically equal to $\lceil \frac{n}{2} \rceil - \log_2 n$ if $|H| = |H'| = n$ (f.g., H and H' consist of one orbit).

Proof. Let $i_1, \dots, i_{\lceil \frac{n}{2} \rceil}$ be arbitrary $\lceil \frac{n}{2} \rceil$ distinct indices, L_b be the dimension $\lceil \frac{n}{2} \rceil$ subspace of F_2^n defined by $x_{i_1} = \dots = x_{i_{\lceil \frac{n}{2} \rceil}} = b$, where $b = 0$ or $b = 1$. We have $S_1(f') \supset S_1(f) - H'$ and $S_1(f')|_{L_0} > 2^{\lceil n/2 \rceil} - 2^d$, where $d = \lceil \log_2 |H| \rceil$. Similarly we have $S_1(1 + f') \supset S_1(1 + f) - H$ and $S_1((1 + f')|_{L_1}) > 2^{\lceil n/2 \rceil} - 2^d$. From Theorem 3, either $AI(f) > \lceil \frac{n}{2} \rceil - \lceil \log_2 |H| \rceil$ or the annihilator functions of f' or $1 + f'$ are zero on L_0 and L_1 . This implies that the monomials in the algebraic normal forms f' and $1 + f'$ have to contain at least $\lceil \frac{n}{2} \rceil$ variables. In the later case $AI(f) = \lceil \frac{n}{2} \rceil$. The conclusion is proved.

IV. Conclusion

We presented a method to obtain some lower bounds on the algebraic immunity for Boolean functions. When the results are applied to symmetric or rotation symmetric Boolean functions, some lower bounds on the algebraic immunity can be proved for these Boolean functions. Some rotation symmetric Boolean functions with their AI near the maximal possible value $\lceil \frac{n}{2} \rceil$ are constructed. Our method suggested some symmetric and rotation symmetric Boolean functions of large number of variables with high algebraic immunity. Thus they can be possibly used in stream ciphers if these Boolean functions satisfy other cryptographic criteria.

Acknowledgement. The work of the 1st author's was supported in part by NNSF of China under Grant 90607005 and Distinguish Young Scholar Grant 10225106.

REFERENCES

- [1] F.Armknecht and M.Krause, Algebraic attacks on stream combiners with memory, in Advances in Cryptology-Crypto2003, LNCS 2729, pages 162-176, Springer-Verlag.
- [2] F.Armknecht, Improving fast algebraic attacks, in Fast Software Encryption -2004, LNCS 3017, pages 65-82, Springer-Verlag.
- [3] F.Armknecht, On the existence of low-degree equations for algebraic attacks, Cryptology e-print Archive, 2004/185
- [4] F.Armknecht, C.Carlet, P.Gaborit, S.Kunzli, W.Meier and O.Ruatta, Efficient computation of algebraic immunity for algebraic and fast algebraic attacks, Advances in Cryptology -Eurocrypt 2006, LNCS 4004, pages 147-164.
- [5] An Braeken and B.Preneel, On the algebraic immunity of symmetric Boolean functions, Indocrypt 2005.
- [6] An Braeken, J.Lano and B.Preneel, Evaluating the resistance of stream ciphers with linear feedback against fast algebraic attacks, ACISP 2006, LNCS 4058, pages 40-51.
- [7] A.Canteaut, Open problems related to algebraic attacks on stream ciphers, In WCC 2005, pages 1-10.
- [8] A.Canteaut and M.Videau, Symmetric Boolean functions, IEEE Transactions on Information theory, vol. 51(2005), no. 8, pages 2791-2811.
- [9] C. Carlet "Boolean Functions for Cryptography and Error Correcting Codes" (150 pages), chapter of the monography "Boolean methods and models" published by Cambridge University Press (Peter Hammer et Yves Crama editors).
- [10] C.Carlet, D.K.Dalai, K.C.Gupta and S.Maitra, Algebraic immunity for cryptographically significant Boolean functions: analysis and construction, IEEE Trans. Inf. Theory, vol.52(2006), no.7, pages 3105-3121.
- [11] C.Carlet, On the Higher Order Nonlinearities of Algebraic Immune Functions, Advances in Cryptology-Crypto 2006, LNCS 4117.
- [12] C.Carlet, A method of construction of balanced functions with optimum algebraic immunity, Cryptology e-print Archive, 2006
- [13] N.Courtois and W.Meier, Algebraic attacks on stream ciphers with linear feedback, in Advances in Cryptology-Eurocrypt 2003, LNCS2656, pages 346-359, Springer-Verlag.
- [14] N.Courtois, Fast algebraic attacks on stream ciphers with linear feedback, in Advances in

Cryptology-Crypto2003, LNCS 2729, pages 176-194, Springer-Verlag.

[15] N.Courtois and J.Pieprzyk, Cryptanalysis of block ciphers with overdetermined systems of equations, in Advances in Cryptology-Asiacrypt2002, LNCS, 2501, pages 267-287.

[16] T.W.Cusick and P. Stanica, Fast evaluation, weighted and nonlinearity of rotation-symmetric functions, Discrete Math., vol.258(2002), pages 289-301.

[17] D.K.Dalai,K.C.Gupta and S.Maitra, Results on algebraic immunity of cryptographically significant Boolean functions, in Indocrypt 2004, LNCS 3348

[18] D.K.Dalai, S.Maitra and S.Sarkar, Basic Theory in construction of Boolean functions with maximal possible annihilator immunity, Cryptology e-print Archive, 2005/229.

[19] J.D.Golic, Vectorial Boolean functions and induced algebraic equations, IEEE Transactions on Information Theory, vol. 52, no. 2, pages 528-537, Feb.2006.

[20] G. Gong, On existence and invariant of algebraic attacks, preprint.

[21] W.Meier, E.Pasalic and C.Carlet, Algebraic attacks and decomposition of Boolean functions, in Advances in Cryptology-Eurocrypt-2004, LNCS 3027, pages 474-491, Springer-Verlag.

[22] J. Pieprzyk and C.X.Qu, Fast hashing and rotation symmetric functions, J.Universal Comput.Sci., vol.5(1999), pages 20-31.

[23] P.Stanica and S.Maitra, A constructive count of rotation symmetric functions, Information processing Letter, vol. 88(2003), pages 299-304.

[24] P.Stanica and S.Maitra, Rotation symmetric Boolean functions -Count and cryptographic properties, preprint.